

Over-expression of a WRKY transcription factor gene *BoWRKY6* enhances resistance to downy mildew in transgenic broccoli plants

Ming Jiang¹ · Qing-e Liu² · Zhen-Ning Liu³ · Jin-Zhi Li¹ · Cai-Ming He¹

Received: 14 March 2016 / Accepted: 27 April 2016
© Australasian Plant Pathology Society Inc. 2016

Abstract WRKY transcription factors play an important role in plant growth, development and immunity. In our study, a WRKY family member gene designated *BoWRKY6* was isolated from broccoli (*Brassica oleracea* var. *italica*), and its expression was induced by downy mildew (*Hyaloperonospora parasitica*). Five transgenic broccoli lines over-expressing *BoWRKY6* driven by the CaMV 35S promoter were obtained by *Agrobacterium tumefaciens* mediated transformation, and they demonstrated significant increased resistance to downy mildew, with resistant levels from low to very high. Real time-qualitative PCR analysis indicated that expressions of both *BoWRKY6* and the pathogenesis-related gene 1 (*BoPRI*) in transgenic plants were obviously higher than those in WT plants after *H. parasitica* treatment. Lines of *BWK14* and *BWK31* exhibited very high resistance to downy mildew, and may serve as promising candidate materials for broccoli molecular breeding in the near future.

Keywords *Brassica oleracea* var. *italica* · Downy mildew · Over-expression · WRKY transcription factor

✉ Ming Jiang
jiangming@tzc.edu.cn

¹ Ecology Key Discipline of Zhejiang Province/College of Life Science, Taizhou University, Jiaojiang 318000, People's Republic of China

² College of Ecology, Lishui University, Lishui 323000, Zhejiang, People's Republic of China

³ Laboratory of Cell and Molecular Biology, Institute of Vegetable Science, Zhejiang University, Hangzhou 310058, People's Republic of China

Introduction

Plants are constantly exposed to a wide range of biotic and abiotic environmental stresses, and extreme temperatures, drought, salinity, chemical toxicity, oxidative stress, bacteria, virus, fungi, nematodes and insect pests are regarded as serious threats to plant production, by affecting vegetative growth, biomass accumulation as well as quality properties (Gupta and Sharma 2013). Abiotic stress factors have a huge impact on crop losses, and it has been estimated that they reduce average yields by more than 50 % for most major crop plants globally (Bray et al. 2000; Wang et al. 2003). Likewise, biotic stresses cause serious impact on crop, and there are approximately 10 000 species of insect pests, 50 000 species of fungal pathogens, 1 800 species of weed plants and 15 000 species of nematodes that destroy agricultural crops (Klassen and Schwartz, 1985; Koul 2011). To cope with these unfavorable environmental conditions, plants have evolved with complex mechanisms, including changes at cellular, molecular, and physiological levels (Akpınar et al. 2012; Atkinson and Urwin 2012). At molecular level, plant stress responses are regulated by diverse signal transduction pathways that are mediated by hormones like salicylic acid (SA), ethylene (ET), jasmonic acid (JA) and abscisic acid (ABA) (Ludwig et al. 2005; Sánchez-Vallet et al. 2012).

A variety of genes have been reported to be induced by environmental stresses, and the induced gene products include late embryogenesis abundant proteins, chaperones, detoxification enzymes, transcription factors, protein kinases as well as phosphoinositide-metabolizing enzymes, which function in protecting cells against oxidative damage (Xiong et al. 2002; Kaur and Gupta 2005). Among them, the cellular abundance of transcription factors are important regulatory proteins that induce or regulate the expression of a series specific genes during stress conditions by interacting with *cis*-elements in

promoter regions, in *Arabidopsis* for example, there are more than 1500 transcription factors covering over 5 % of its genome, and AP2/EREBP, ABI3/VP1, ARF, bHLH, bZIP, HB, HSF, MYB, NAC as well as WRKY are recognized as known stress responsive transcription factor families (Riechmann et al. 2000; Shameer et al. 2009).

WRKY domain has been defined as an approximately 60 amino acid motif sequence of WRKYGQK at its N-terminus as well as a zinc finger structure of C2H2 or C2HC (C and H indicate cysteine and histidine amino acid residues, respectively.) at C-terminus (Eulgem et al. 2000; Zhang and Wang 2005). The WRKY gene family participates in plant growth, development, and stress responses. Though discovered relatively recently, they have been regarded as one of the best characterized transcription factor families (Chen et al. 2012). However, to our knowledge, little is known about the biological functions of WRKYs in downy mildew (*Hyaloperonospora parasitica*) resistance. Downy mildew is one of the most destructive diseases of cruciferous crops, and it causes serious problems in commercial production of cabbage, broccoli, cauliflower, radish and mustard. Broccoli (*Brassica oleracea* var. *italica*) is susceptible to infection by downy mildew at all growth stages, causing damages on leaves, stems as well as flower parts, which consequently decreased flower head yield and quality (Wang et al. 2001). In our study, a *Brassica oleracea* var. *italica* WRKY gene, designated *BoWRKY6*, was isolated and introduced into broccoli driven by the CaMV 35S promoter, and five broccoli lines illustrating significantly higher resistance against downy mildew were obtained.

Materials and methods

Plant and fungal materials

Broccoli (*B. oleracea* var. *italica*) line Bo112 (highly resistant to downy mildew) was cultured in a growth chamber to two-leaf stage at 25 °C with a photoperiod of 16 h/8 h. Downy mildew infected broccoli leaves were collected in our test field in Linhai, China. Conidial cell suspensions were prepared by rinsing the conidia off the infected leaves, and 0.2 mL of suspension (approximately 1 000 conidia mL⁻¹) were sprayed onto each side of true leaves, and the control leaves were applied with an equal amount of sterile water. The seedlings were sealed to maintain an approximately 85–100 % relative humidity, and then cultured in a growth cabinet at 16 °C under a 16 h/8 h light/dark cycle. Leave samples were collected at 0, 6, 12, 24, 36 and 72 h after inoculation, and stored at - 80 °C for RNA extraction.

Isolation of *BoWRKY6* gene

Genomic DNA was extracted from broccoli leaf samples following cetyl trimethyl ammonium bromide (CTAB) protocol

(Doyle and Doyle, 1987). Total RNA of each treatment was isolated with Trizol reagent (Invitrogen, USA), and cDNA was synthesized using the SMART cDNA PCR KIT (Clontech, USA) according to its manual. *BoWRKY6* was amplified by a primer pair with restriction enzyme sites (WK6UP1: 5'-GCATCCCGGGATGGATATTGCTAGTAATAACAA-3'; WK6DN1: 5'-ATATGAGCTCAACCGAAACTGCTGGC-3'). PCR reactions were performed in 20 µL volume containing 30 ng template DNA or cDNA, 25 pmol of each primer, 0.8 U of DNA polymerase (Promega, UK), 0.45 µM dNTP mixture, 50 mM Tris-HCl (pH 8.3), and 2.5 mM MgCl₂. The *BoWRKY6* was amplified with the following cycling profile: 95 °C for 5 min, followed by 32 cycles at 95 °C for 30 s, 55.8 °C for 40 s, and 72 °C for 80 s, and a final extension at 72 °C for 10 min. PCR products were electrophoresed on a 1.2 % agarose gel containing ethidium bromide. Bands of interest were excised, and then purified by using QIAquick Gel Extraction Kit (QIAGEN, Germany). The purified PCR products were cloned into pGEM-T EASY vectors (Promega, UK), and *Escherichia coli* JM109 was then transformed with the recombinant plasmids. Five positive clones were selected by blue/white screening on Luria-Bertani (LB) plates containing 100 µg/mL ampicillin, and were then sequenced.

Sequence analysis

Homologous protein sequences were downloaded from NCBI (www.ncbi.nlm.nih.gov) for sequence comparisons and phylogenetic analysis. Those data included *B. napus* (ACQ76810.1), *B. rapa* (AHB33862.1), *Arabidopsis lyrata* subsp. *lyrata* (XP_002878082.1), *Eutrema salsugineum* (XP_006402991.1), *A. thaliana* (NP_191199.1), *Thellungiella halophila* (BAJ33964.1), *Capsella rubella* (XP_006295843.1), *Jatropha curcas* (AGQ04251.1), *Populus trichocarpa* (XP_002319879.2), *Theobroma cacao* (XP_007011367.1), *Citrus clementina* (XP_006435943.1), *Ricinus communis* (XP_002520871.1), *Phaseolus vulgaris* (XP_007136415.1), *Glycine max* (XP_003526799.1), *Cicer arietinum* (XP_004502820.1), *Brachypodium distachyon* (XP_003566949.1), and *Oryza sativa* (AAU44093.1). Multiple alignments of *BoWRKY6* and its homologous protein sequences were developed with ClusterX 1.81 (Thompson et al. 1997), and phylogenetic trees were constructed by Neighbor-joining method using Molecular Evolutionary Genetic Analysis (MEGA) 3.1 version with 1 000 bootstrap replications (Kumar et al. 2004).

Expression analysis of *BoWRKY6* gene

According to sequencing results, primer pairs WK6UP2 (5'-GATTCCTTCGAACCCATCTC-3')/WK6DN2 (5'-TCAGAATCCGTGATGATCTCATC-3') were designed and

employed for expression analysis. RT-PCR assays were conducted using 30 ng cDNA templates of downy mildew treated and control leaves, employing the same cycling conditions as gene cloning. PCR products were separated on a 1.0 % agarose gel, and then photographed using Gel Doc XR⁺ System (BIO-Rad, UK). Actin gene was used as an internal control, and two primers, ACTUP (5'-TCTCGATGGAAGAGC TGGTT-3') and ACTDN (5'-GATCCTTACCGAGGGA GGTT-3'), were applied to amplify actin gene fragments using following profile: 94 °C for 5 min; 32 cycles of 94 °C for 30 s, 55.6 °C for 45 s, and 72 °C for 60 s; followed by a final extension at 72 °C for 7 min.

Expression vector construction and transformation

Purified PCR products were digested with two restriction enzymes of *Sma* I and *Sac* I, and then cloned into binary vector pBI121 vector which was digested by the same two enzymes. The recombinant vector was then introduced into *Agrobacterium tumefaciens* strain LBA4404 competent cells. Broccoli line Bo0283 (susceptible to downy mildew) was used for transgenic study. Stems of 18-day old seedlings were harvested, and dipped in 0.1 % HgCl₂ for 9 min, and were then inoculated with *A. tumefaciens*. Murashige and Skoog (MS) supplemented with 0.02 mg/L of α -naphthaleneacetic acid (NAA), 4.0 mg/L of 6-benzylaminopurine (6-BA) and 5.0 mg/L of AgNO₃ was prepared for pre- and co-culture mediums. Shoot induction medium was made up of MS plus 0.02 mg/L of NAA, 4.0 mg/L of 6-BA, 4.0 mg/L of AgNO₃ and 50.0 mg/L of kanamycin (Km). Regenerated shoots were rooted on half strength MS medium containing 0.2 mg/L of NAA and 50.0 mg/L of Km (Jiang et al. 2012).

Screening of transgenic plants

Genome DNA of broccoli transgenic lines was isolated using CTAB method, and primer pair WK6UP3 (5'-TCAACA AAGGGTAATATCCGG-3')/WK6DN3 (5'-CAAGGCTT GATTTTGGGTG-3') was employed to amplify fragments with partial 35S promoter and *BoWRKY6* gene. Approximately 30 ng DNA of each line was used as PCR templates. The PCR conditions were as follows: 95 °C for 5 min; 32 cycles of 95 °C for 30 s, 53.5 °C for 50 s, and 72 °C for 75 s; followed by a final extension at 72 °C for 10 min. PCR products were electrophoretically separated on a 1.0 % agarose gel at 120 V for 30 min, and then photographed using the Gel Doc XR⁺ System (BIO-Rad, UK).

Expression of *BoWRKY6* and pathogenesis-related gene 1

WT and transgenic plants were inoculated with *H. parasitica*, and leaf samples were collected at 0 h and 24 h after infection.

Total RNA was isolated from 100 mg of leaves using Trizol (Invitrogen, USA), and the 1st cDNA was synthesized using the SMART cDNA PCR KIT (Clontech, USA). For Real time-qualitative PCR (RT-qPCR), three primer pairs were used to determine gene expression of *BoWRKY6*, *BoPRI* (pathogenesis related gene 1), and *BoActin* (a reference gene) (Lovelock et al. 2013). The sequences of gene specific primers were as follows: BoWRKY6rtup (TATCTCGTCACAGC CGCC), BoWRKY6rtdn (AACTGGTCCCAATCTTTTCT), BoPRIup (GCGACTGCAGACTCGTACAC), BoPRI1dn (TCTCGTTGACCCAAAGGT), BoActinINup (ACGTGGACATCAGGAAGGAC) and BoActindn (GAACCACCGATCCAGACACT). All the qRT-PCR runs were carried out in a LightCycler[®] 96 real-time PCR System (Roche, Switzerland) by using YBR FastStart Essential DNA Green Master Mix (Roche, Switzerland). The following reagents were added to 20 μ l tubes: 10 μ l of Master Mix; 0.2 μ l of each primer (20 μ M), 3 μ l of diluted cDNA(0 h and 24 h), and 6.6 μ l of ddH₂O. The following PCR conditions were used: 95 °C for 10 min, followed by 40 cycles of 95 °C for 15 s; 55 °C for 15 s, and 72 °C for 30 s. 60 °C for 1 min; 72 °C for 30 s. All qRT-PCR assays were conducted in triplicate. Mean fold changes in gene expression of *BoWRKY6* and *BoPRI* were normalized against *BoActin* gene by using the equation as described by Livak and Schmittgen (2001).

Disease resistance assessment and statistical analysis

Interaction phenotypes were determined 7 d after inoculation. Scales ranging from 0 to 9 were used for classifying leaf infection by downy mildew pathogen: 0 = no necrotic flecks, no sporulation; 1 = small necrotic flecks, no sporulation; 3 = necrotic flecks, one to few sporangiophores; 5 = necrotic lesions, sparse scattered sporulation usually confined to necrotic areas; 7 = necrotic lesions, sometimes with accompanying chlorosis, scattered, heavy to abundant sporulation in both chlorotic and necrotic areas; 9 = necrosis and some chlorosis may be evident, uniformly heavy sporulation over abaxial surface of leaf (Li et al. 2010). The data were statistically analyzed by using one-way analysis of variance (ANOVA) and Duncan's test at a significance level of $p=0.05$.

Results

Isolation and characterization of *BoWRKY6*

Primer pair WK6UP1/WK6DN1 was employed to amplify *BoWRKY6* gene by using leaf genome DNA and cDNA as templates, respectively. Sequencing results indicated *BoWRKY6* was 1498 bp in length, with two introns of 467 and 179 bp (Fig. 1). The complete coding sequence of *BoWRKY6* was 852 bp in length encoding 283 amino acids,




Fig. 1 Gene structure of *BoWRKY6*. Boxes indicate exons; black lines represent introns

carrying a nuclear localization sequence (NLS) of PVKGRKRCYKRKKK, and a WRKY motif of WRKYGQK, together with a zinc finger of C-X₇-C-X₂₃-H-X₁-C (X represents any amino acid) at its C-terminal (Fig. 2).

Sequence comparison and phylogenetic analysis of *BoWRKY6*

For sequence comparison and phylogenetic analysis, 17 homologous protein sequences were downloaded from NCBI, and were aligned with ClusterX 1.81. These protein sequences together with *BoWRKY6* varied in length, ranging from 276 (*B. rapa*) to 338 (*T. halophila*), with an average size of 299. *BoWRKY6* demonstrated the highest degree of homology to *B. napus* (CDY32013.1) and *B. rapa* (NP_001288847.1) by performing BLAST searches via NCBI, with identities of 99 and 94 %, respectively, and the lowest homology were observed in *B. distachyon* (XP_003566949.1) and *O. sativa* (AAU44093.1), with sequence identities less than 40 %. However, the 18 WRKY domains were relatively well conserved both in length and amino acid composition, and each domain contained a WRKYGQK motif as well as a C-X₇-C-X₂₃-H-X₁-C pattern zinc finger sequence (Fig. 3).

To understand the relationship between *BoWRKY6* and its homologous sequences, phylogenetic tree was constructed

using Mega software. Based on the results, 18 WRKY proteins were divided into 4 distinct groups, namely I, II, III and IV (Fig. 4). It could be observed that *BoWRKY6* shared a close lineage with other Cruciferae plants of *B. napus*, *B. rapa*, *E. salsugineum*, *T. halophila*, and *C. rubella*, with 100 % bootstrap confidence value. WRKY proteins of three Leguminosae plants, *C. arietinum*, *P. vulgaris* and *G. max*, were closely related in our phylogenetic tree, with confidence of 100 %. WRKY sequences of *J. curcas*, *C. clementina*, *R. communis*, *P. trichocarpa* and *T. cacao* group together, however, the confidence value was lower (83 %) when compared to group I, II, and IV. Group IV was consisted of two Gramineae species, *B. distachyon* and *O. sativa*, with 100 % bootstrap confidence value.

Expression analysis

RT-PCR was performed to determine the expression patterns of *BoWRKY6* in both control and downy mildew challenged leaves. Results indicated that *BoWRKY6* expressed constitutively in control leaves at a relatively lower level, however, increasing levels of expression were detected 6–24 h after inoculation, and then decreased at 36 h (Fig. 5).

Screening of transgenic plants

BoWRKY6 gene driven by a cauliflower mosaic virus (CaMV) 35S promoter was transformed into broccoli, and T₀ plants with abnormal morphological characteristics were discarded. Totally five transgenic lines, namely *BWK05*, *BWK14*, *BWK16*, *BWK31* and *BWK54*, which exhibited normal growth and developmental traits, were screened out of 75 kanamycin-resistant plants, and they were later confirmed by PCR using

```

1   ATGGATATTGCTAGTAATAACAAGCAATAAAGCTAAAAGTTAGGGACCAACTTCTTCAATGCCACGAGATGACCACTAAG
1   M D I A S N N K A I K L K V R D Q L L Q C H E M T T K
82  GTTCAGCAACTCCTCTCTCAAGACGGGTTCGGATTGGGTCCAGCGAAGGATCTCGTGGAGAAAATATTGGGGTCTATCAGT
28  V Q Q L L S Q D G S D L G P A K D L V E K I L G S I S
163 GACAAATCTCTGCTTGTGATTCTTCGAACCCATCTCCCCCTCTTATCTCGTCACAGCCGCCGAAGGCTCTCAAAATGCT
55  D T I S A L D S F E P I S P S Y L V T A A E G S Q N A
244 TCCTGCGACAACGACGGCAAGCTTGAGGATTCTGGCGATAGTCAGAAAAGATTGGGACCAAGTTAAGGGTAAAAGAGGATGC
82  S C D N D G K L E D S G D S Q K R L G P V K G K R G C
325 TACAAGAGAAAAGAAGAAATCAGAGACGTGGACTGTAGAGTCTACCGTACTTGAGGACACATTTTCTTGGAGGAAATATGGA
109  Y K R K K K S E T W T V E S T V L E D T F S W R K Y G
406 CAAAAACAGATTCTTAATGCCAAATTCCTCAAGAAGTTACTTTAGGTGCACACACAAATACACTCAAGGGTCAAGGCAACA
136  Q K Q I L N A K F P R S Y F R C T H K Y T Q G C K A T
487 AAGCAAGTGCAGAAGCTAGAGTCTGAACCCAGGATGTTTCAGCATCACATACATCGGAAACCCACAGTGTAAATACCAACGAA
163  K Q V Q K L E S E P R M F S I T Y I G N H T C N T N E
568 GTAACACCCAAAATCAAGCCTTGTATTTCATCATGATGAGATCATCAGGATTTCTGAAGAGATCCAGAGTCTAGTTTGTATG
190  V T P K I K P C I H H D E I I T D S E E I Q S P S L M
649 ACCTCGATGAAGGAAGAGGAAGAAAATCACCATCATGGTTTCGTCACAGGAGTACTGACTTGGTGTGGTGTGGTGTGGTGTGG
217  T S M K E E E E N H H H G S S T E S D L Q L V W Q E M
730 TTGGTCTTTGACAGGAAACACCATCATCATCAGGCTGTTTACGGTGTGGGGAACTAGTACATCTATCAATGGTTTG
244  L V F A E E H H H H E A V Y G C G E T S T S I N G L
811 GATTCGCGGATCTGTGGAGTTGCCAGCAGTTTTTCGGTTTTAG
271  D S A D L W S C Q Q F S V *

```

Fig. 2 Complete coding sequence of *BoWRKY6* and its deduced amino acids. Dashed line indicates a nuclear localization sequence; Line represents the WRKY motif; Boxed sequence indicates a zinc finger domain




Fig. 3 Comparisons of WRKY domains between BoWRKY6 and its homologous sequences

WK6UP3/WK6DN3 primer pair. Fragments containing partial 35S promoter and *BoWRKY6* were amplified only in transgenic lines, and no band was observed in WT line (Fig. 6).

Expression analysis of BoWRKY6 and BoPR1

There was no obvious difference in the basal expression levels of *BoWRKY6* between the WT and transgenic leaves. However, *BoWRKY6* transcripts increased greatly after infiltration with *H. parasitica*, and higher expression levels at 24 h after infection were observed in all transgenic lines when compared to WT. The expression levels increased 2.11 times in WT, while in transgenic plants, 3.44 to 9.85 times of expression levels were detected when infected by *H. parasitica* in 24 h (Fig. 7).


PR1 is regarded as a marker gene of the SA signaling pathway, and in our present study, qRT-PCR was carried out

to analyze its expression level in both WT and transgenic broccoli lines. The results indicated no significant difference before *H. parasitica* infection (0 h), however, after 24 h of *H. parasitica* inoculation, *BoPR1* expression levels increased 2.15 times in WT plants, and it increased 6.62 to 13.90 times in those of transgenic lines (Fig. 7).

Disease resistance evaluation

To evaluate the resistance of five transgenic lines to downy mildew, enough T₀ broccoli plants were produced using tissue culture method, and 30 plants per line were assessed, with three replicates. Disease reaction phenotypes were made at 7 d post-inoculation, and disease indices were calculated (Table 1). Significant differences in disease index were observed between WT and transgenic lines, and the WT plants demonstrated a susceptible reaction with disease index of

Fig. 4 Phylogenetic tree of BoWRKY6 and its homologous sequences using Neighbor-joining method






Fig. 5 Expression patterns of *BoWRKY6* gene in response to downy mildew inoculation. **a**: Leaves challenged by downy mildew; **c**: control leaves; **b** and **d**: internal control of actin gene

6.97, while the *BoWRKY6* over-expressing lines showed different levels of downy mildew resistance, with resistance classifications ranging from LR to VR. Necrotic lesions, chlorosis, and heavy sporulation were observed on WT leaves, whereas lines *BWK16* and *BWK54* exhibited fewer necrotic lesions, less chlorosis and light sporulation. *BWK31* and *BWK14*, which were identified as very resistant (VR), exhibited only small necrotic flecks and few sporangiophores on infected leaves.

Discussion

WRKY transcription factors are regarded as one of the largest families of transcriptional regulators in higher plants, and they form sophisticated signaling networks that modulate multiple plant processes (Rushton et al. 2010). Owing to the availability of increasing numbers of sequenced genomes, WRKY gene families were identified and characterized within many plants. In *Klebsormidium flaccidum*, a charophyte plant, only two WRKY genes are present in its genome, and they belong to Group I and IIb, respectively (Rinerson et al. 2015). However, the gene numbers expanded due to plant evolution, which is likely to be associated with the generation of highly complex defence mechanisms (Rushton et al. 2010). In the model plant of *Arabidopsis*, there are 72 WRKY genes, and 49 of them are differentially regulated when challenged by an avirulent strain of *Pseudomonas syringae* or treated by salicylic acid (SA) (Dong et al. 2003). Whereas in *Brachypodium distachyon*, 86 WRKY genes were identified, and dozens of *BdWRKYs* were rapidly and significantly up-regulated after inoculation of *Pseudomonas syringae* or *Boea hygrometrica* (Wen et al. 2014). There are more WRKY genes in *Oryza sativa* than in *Arabidopsis thaliana* and *B. distachyon*, respectively 98 and 102 WRKY genes were identified in *japonica* as well as in *indica* rice (Ross et al. 2007).

WRKYGQK is a common signature motif present in WRKY transcription factors, and is directly involved in DNA binding, with W-box element (C/T)TGAC(C/T) in the promoter region as its target (Eulgem and Somssich 2007).

WT *BWK05* *BWK14* *BWK16* *BWK31* *BWK54*



Fig. 6 PCR confirmation of *BoWRKY6* over-expressing broccoli plants

Though WRKYGQK is highly conserved in most WRKY members, variant proteins such as WRKYGKK, WRKYGEK, WRKYEDK, WKKYGQK, CRKYGQK, WHQYGLK, WRKYGMK, WSKYGQK, WQKYGQK as well as WIKYGEN were identified in *Solanum lycopersicum*, *Glycine max* and *Vitis vinifera* (Huang et al. 2012; Bencke-Malato et al. 2014; Guo et al. 2014). WRKY motif sequence is followed by a C2H2- or C2HC-type zinc finger motif, with sequences of C-X₄₋₅-C-X₂₂₋₂₃-H-X-H and C-X₅₋₈-C-X₂₅₋₂₈-H-X₁₋₂-C, however, there are some exceptions, e.g. in *Salvia miltiorrhiza*, six out of 61 WRKY members contain a C2HC zinc finger motif of C-X₇-C₂₃-H-X₁-C. In our study, a WRKY transcription factor gene was isolated from broccoli, with a WRKY motif of WRKYGQK and a zinc finger of C-X₇-C-X₂₃-H-X₁-C.

Plants are often subjected to abiotic and biotic stresses in their natural habitat, and adapt to such changes requires some degree of phenotypic plasticity that is mainly determined by its genome (Pandey and Somssich 2009). WRKY transcription factors play important roles not only in plant growth and development, but also in adaptive plasticity of the highly variable environments. WRKY transcription factors participate in plant immunity by regulating genes with W box elements within their promoters or by interacting with other transcription factors (Pandey and Somssich 2009). Expression of *GhWRKY15* was significantly induced in *Gossypium hirsutum* seedlings when treated with fungal, salicylic acid (SA), methyl jasmonate or methyl viologen, and tobacco over-expressing *GhWRKY15* demonstrated more resistance to *Colletotrichum*




Fig. 7 Expression of *BoWRKY6* and *BoPRI* genes in WT and transgenic broccoli leaves in response to downy mildew by using qRT-PCR. Each column represents the average of three replicates, and error bars represent the standard deviation

Table 1 Evaluation of resistance to downy mildew in transgenic lines

Lines	Interaction-phenotype class						Plant assayed	Disease index ^a	Resistance classification ^b
	0	1	3	5	7	9			
CK				41	40	39	120	6.97 a	S
BWK05		34	47	39			120	3.08 d	MR
BWK14	23	33	37	27			120	2.33 e	VR
BWK16			33	43	44		120	5.18 b	LR
BWK31	31	42	42	5			120	1.61 f	VR
BWK54			38	40	37		120	4.78 c	LR

^a Means within each column followed by the same letter are not significantly different ($p < 0.05$)

^b Resistance classes based on disease indices (DI) calculated by Williams' formula: VR (very resistant), DI = 0–3.0; MR (moderately resistant), DI = 3.1–5.0; LR (low resistance), DI = 5.1–6.0; S (susceptible), DI = 6.1–7.0; VS (very susceptible), DI = 7.1–9.0

gossypii and *Phytophthora parasitica* (Yu et al. 2012). *S. lycopersicum* defense-related WRKY1 gene *SIDRW1* was significantly induced by *Botrytis cinerea*, and silencing of this gene resulted in increased severity of disease caused by *B. cinerea* (Liu et al. 2014). A *Populus trichocarpa* WRKY gene named *PtrWRKY73* was induced by exogenous SA, its over-expression in *Arabidopsis thaliana* resulted in increased resistance to *Pseudomonas syringae* (PstDC3000) (Duan et al. 2015). In present study, a WRKY gene designated *BoWRKY6* was isolated from broccoli, and its expression was induced by downy mildew. Five transgenic lines over-expressing *BoWRKY6* demonstrated increased resistance to downy mildew, with resistance classification from low to very high resistant.

In plants, disease resistance is regulated by multiple signal transduction pathways, and phytohormones such as salicylic acid (SA), jasmonic acid (JA) and ethylene (ET) act as key signaling molecules which induce different transcriptional regulation of pathogenesis-related (PR) genes (Maleck et al. 2002; Lee et al. 2014). SA pathway play an important role in defense response to biotrophic pathogens, and *PR1*, a reliable molecular marker for systemic acquired resistance, is dependent on SA perception and associated with induced resistance against the biotrophic fungus *H. parasitica* in *A. thaliana* (Thomma et al. 1998; Maleck et al. 2000). Over-expression of an *Arabidopsis* cysteine-rich receptor-like kinase gene *CRK13* exhibited resistance to *Pseudomonas syringae*, and high level transcript accumulations of *PR1* and other PR genes were observed (Acharya et al. 2007). In our current study, transgenic broccoli plants over-expressing the *BoWRKY6* gene demonstrated increased expression of *PR1*, which is correlated positively with enhancement of downy mildew resistance in those lines.

In conclusion, a WRKY gene namely *BoWRKY6* was isolated from broccoli, with complete coding sequence of 852 bp in length encoding 283 amino acids, carrying a WRKY motif of WRKYGQK and a zinc finger of C-X₇-C-X₂₃-H-X₁-C.

Expression of *BoWRKY6* was induced by downy mildew, and transgenic plants over-expressing *BoWRKY6* showed significant resistance to downy mildew, with resistant levels from low to very high. Two lines of *BWK14* and *BWK31* demonstrated very high resistance to downy mildew, and may serve as promising candidate breeding materials in the future.

Acknowledgments This work was supported by Zhejiang Provincial Natural Science Foundation of China (LY13C150003), Public Welfare Technology Research Projects of Zhejiang Province (2016C32091), Ecology Key Disciplines of Zhejiang Province in Taizhou University (EKD 2013-03).

References

- Acharya BR, Raina S, Maqbool SB, Jagadeeswaran G, Mosher SL, Appel HM, Schultz JC, Klessig DF, Raina R (2007) Overexpression of CRK13, an *Arabidopsis* cysteine-rich receptor-like kinase, results in enhanced resistance to *Pseudomonas syringae*. *Plant J* 2007(50):488–499
- Akpinar BA, Avsar B, Lucas SJ, Budak H (2012) Plant abiotic stress signaling. *Plant Signal Behav* 7:1450–1455
- Atkinson NJ, Urwin PE (2012) The interaction of plant biotic and abiotic stresses: from genes to the field. *J Exp Bot* 63:3523–3543
- Bencke-Malato M, Cabreira C, Wiebke-Strohm B, Bücken-Neto L, Mancini E, Osorio MB, Homrich MS, Turchetto-Zolet AC, De Carvalho MC, Stolf R, Weber RL, Westergaard G, Castagnaro AP, Abdelnoor RV, Marcelino-Guimarães FC, Margis-Pinheiro M, Bodanese-Zanettini MH (2014) Genome-wide annotation of the soybean WRKY family and functional characterization of genes involved in response to *Phakopsora pachyrhizi* infection. *BMC Plant Biol* 14:236
- Bray EA, Bailey-Serres J, Weretilnyk E (2000) Responses to abiotic stresses. In: Gruissem W, Buchanan B, Jones R (eds) *Biochemistry and molecular biology of plants*. American Society of Plant Physiologists Press, Rockville, pp 1158–1249
- Chen L, Song Y, Li S, Zhang L, Zou C, Yu D (2012) The role of WRKY transcription factors in plant abiotic stresses. *Biochim Biophys Acta* 1819:120–128
- Dong J, Chen C, Chen Z (2003) Expression profiles of the *Arabidopsis* WRKY gene superfamily during plant defense response. *Plant Mol Biol* 51:21–37

- Doyle JJ, Doyle JL (1987) A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem Bull* 19:11–15
- Duan Y, Jiang Y, Ye S, Karim A, Ling Z, He Y, Yang S, Luo K (2015) PtrWRKY73, a salicylic acid-inducible poplar WRKY transcription factor, is involved in disease resistance in *Arabidopsis thaliana*. *Plant Cell Rep* 34:831–841
- Eulgem T, Rushton PJ, Robatzek S, Somssich IE (2000) The WRKY superfamily of plant transcription factors. *Trends Plant Sci* 5:199–206
- Eulgem T, Somssich IE (2007) Networks of WRKY transcription factors in defense signaling. *Curr Opin Plant Biol* 10:366–371
- Guo C, Guo R, Xu X, Gao M, Li X, Song J, Zheng Y, Wang X (2014) Evolution and expression analysis of the grape (*Vitis vinifera* L.) WRKY gene family. *J Exp Bot* 65:1513–1528
- Gupta OP, Sharma P (2013) Potential role of small RNAs during stress in plants. In: Gaur RK, Sharma P (eds) *Molecular approaches in plant abiotic stress*. CRC Press, Boca Raton, pp 67–89
- Huang S, Gao Y, Liu J, Peng X, Niu X, Fei Z, Cao S, Liu Y (2012) Genome-wide analysis of WRKY transcription factors in *Solanum lycopersicum*. *Mol Genet Genomics* 287:495–513
- Jiang M, He CM, Miao LX, Zhang YC (2012) Overexpression of a broccoli defensin gene BoDFN enhances downy mildew resistance. *J Integr Agric* 11:1137–1144
- Kaur N, Gupta AK (2005) Signal transduction pathways under abiotic stresses in plants. *Curr Sci* 88:1771–1780
- Klassen W, Schwartz PH (1985) ARS research program in chemical insect control. In: Hilton JL (ed) *Agricultural chemicals of future*. Rowman and Allanheld, New Jersey, pp 267–291
- Koul O (2011) Microbial biopesticides: opportunities and challenges. *CAB Rev Perspect Agric Vet Sci Nutr Nat Resour* 6:1–26
- Kumar S, Tamura K, Nei M (2004) MEGA3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinform* 5:150–163
- Lee YH, Kim SH, Yun BW, Hong JK (2014) Altered cultivar resistance of kimchi cabbage seedlings mediated by salicylic acid, jasmonic acid and ethylene. *Plant Pathol J* 2014(30):323–329
- Li X, Xia B, Jiang Y, Wu Q, Wang C, He L, Peng F, Wang R (2010) A new pathogenesis-related protein, LrPR4, from *Lycoris radiata*, and its antifungal activity against *Magnaporthe grisea*. *Mol Biol Rep* 37:995–1001
- Liu B, Hong YB, Zhang YF, Li XH, Huang L, Zhang HJ, Li DY, Song FM (2014) Tomato WRKY transcriptional factor SIDRW1 is required for disease resistance against *Botrytis cinerea* and tolerance to oxidative stress. *Plant Sci* 227:145–156
- Livak KJ, Schmittgen TG (2001) Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta C_T}$ method. *Methods* 25:402–408
- Lovelock DA, Donald CE, Conlan XA, Cahill DM (2013) Salicylic acid suppression of clubroot in broccoli (*Brassica oleracea* var. *italica*) caused by the obligate biotroph *Plasmodiophora brassicae*. *Aust Plant Pathol* 42:141–153
- Ludwig AA, Saitoh H, Felix G, Freymark G, Miersch O, Wasternack C, Boller T, Jones JD, Romeis T (2005) Ethylene-mediated cross-talk between calcium-dependent protein kinase and MAPK signaling controls stress responses in plants. *Proc Natl Acad Sci U S A* 102:10736–10741
- Maleck K, Levine A, Eulgem T, Morgan A, Schmid J, Lawton KA, Dang JL, Dietrich RA (2000) The transcriptome of *Arabidopsis thaliana* during systemic acquired resistance. *Nat Genet* 26:403–410
- Maleck K, Neuenschwander U, Cade RM, Dietrich RA, Dangl JL, Ryals JA (2002) Isolation and characterization of broad-spectrum disease-resistant *Arabidopsis* mutants. *Genetics* 160:1661–1671
- Pandey SP, Somssich IE (2009) The role of WRKY transcription factors in plant immunity. *Plant Physiol* 150:1648–1655
- Riechmann JL, Heard J, Martin G, Reuber L, Jiang C, Keddie J, Adam L, Pineda O, Ratcliffe OJ, Samaha RR, Creelman R, Pilgrim M, Broun P, Zhang JZ, Ghandehari D, Sherman BK, Yu G (2000) *Arabidopsis* transcription factors: genome-wide comparative analysis among eukaryotes. *Science* 290:2105–2110
- Rinerson C, Rabara RC, Tripathi P, Shen QJ, Rushton PJ (2015) The evolution of WRKY transcription factors. *BMC Plant Biol* 15:66
- Ross CA, Liu Y, Shen QJ (2007) The WRKY gene family in rice (*Oryza sativa*). *J Integr Plant Biol* 49:827–842
- Rushton PJ, Somssich IE, Ringler P, Shen QJ (2010) WRKY transcription factors. *Trends Plant Sci* 15:247–258
- Sánchez-Vallet A, López G, Ramos B, Delgado-Cerezo M, Riviere MP, Llorente F, Fernández PV, Miedes E, Estevez JM, Grant M, Molina A (2012) Disruption of abscisic acid signaling constitutively activates *Arabidopsis* resistance to the necrotrophic fungus *Plectosphaerella cucumerina*. *Plant Physiol* 160:2109–2124
- Shameer K, Ambika S, Varghese SM, Karaba N, Udayakumar M, Sowdhamini R (2009) STIFDB-*Arabidopsis* stress responsive transcription factor database. *Int J Plant Genomics* 2009:1–8
- Thomma BPHJ, Eggermont K, Penninckx IAMA, Mauch-Mani B, Vogelsang R, Cammue BPA, Broekaert WF (1998) Separate jasmonate-dependent and salicylate-dependent defense-response pathways in *Arabidopsis* are essential for resistance to distinct microbial pathogens. *Proc Natl Acad Sci U S A* 95:15107–15111
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* 25:4876–4882
- Wang M, Farnham MW, Thomas CE (2001) Inheritance of true leaf stage downy mildew resistance in broccoli. *J Am Soc Hortic Sci* 126:727–729
- Wang WX, Vinocur B, Altman A (2003) Plant responses to drought, salinity and extreme temperatures: towards genetic engineering for stress tolerance. *Planta* 218:1–14
- Wen F, Zhu H, Li P, Jiang M, Mao W, Ong C, Chu Z (2014) Genome-wide evolutionary characterization and expression analyses of WRKY family genes in *Brachypodium distachyon*. *DNA Res* 21:327–339
- Xiong L, Schumaker KS, Zhu JK (2002) Cell signaling during cold, drought and salt stress. *Plant Cell* 14:S165–S183
- Yu F, Huaxia Y, Lu W, Wu C, Cao X, Guo X (2012) GhWRKY15, a member of the WRKY transcription factor family identified from cotton (*Gossypium hirsutum* L.), is involved in disease resistance and plant development. *BMC Plant Biol* 12:144
- Zhang Y, Wang L (2005) The WRKY transcription factor superfamily: its origin in eukaryotes and expansion in plants. *BMC Evol Biol* 5:1