



Expression of genes involved in the anthocyanin biosynthesis pathway in white and red fruits of *Fragaria pentaphylla* and genetic variation in the dihydroflavonol-4-reductase gene



Wenkai Duan^{a, b, c}, Peilong Sun^a, Junmin Li^{b, *}

^a Laboratory of Food Energy, Zhejiang University of Technology, Hangzhou 310014, PR China

^b Zhejiang Provincial Key Laboratory of Plant Evolutionary Ecology and Conservation, Taizhou University, Taizhou 318000, PR China

^c Taizhou Vocational College of Science and Technology, Taizhou 318000, PR China

ARTICLE INFO

Article history:

Received 6 March 2017

Received in revised form

20 April 2017

Accepted 22 April 2017

Keywords:

Anthocyanin

Real-time quantitative PCR

Single nucleotide polymorphisms

Wild strawberry

ABSTRACT

Structural and regulatory genes control fruit colors in plants. Real-time quantitative PCR results showed significantly higher expression levels of structural genes (*FpCHS*, *FpDFR*, *FpANS*, and *FpUGFT*) as well as of the regulatory gene *MYB10* in red fruits of *Fragaria pentaphylla* compared to white fruits. These genes were strongly associated with anthocyanin accumulation within fruits. The full-length sequence of the *FpDFR* gene in red fruits of *F. pentaphylla* had a length of 2080 bp, was separated by five introns, and shared 95% homology with the *F. vesca* *DFR* sequence. Twenty-seven SNPs were detected in the *FpDFR* gDNA sequences between red and white fruits. Among these, transition substitutions were more frequent than transversions (66.7% vs. 33.3%), and a larger number of nucleotide variants existed in introns compared to exons (70.4% vs. 29.6%). A Chi-square test showed only three SNPs significantly associated with fruit color. Combined with structural analyses of the *FpDFR* protein and an expression analysis of the anthocyanin pathway genes, these results indicate that *trans*-regulation might contribute to color control in *F. pentaphylla*.

© 2017 Published by Elsevier Ltd.

1. Introduction

Fruit color results from the accumulation of anthocyanin and is an important quality trait, influencing human consumption (Goto and Kondo, 1991). Recently, anthocyanins have attracted increasing interest due to their important physiological functions, such as antioxidative (Wang et al., 1999; Kähkönen and Heinonen, 2003), antimutagenic (Gasiorowski et al., 1997), and anticancer activities (Koide et al., 1996).

The biosynthetic pathway of anthocyanins has been well-described in various plants, including *Zea mays* L., *Antirrhinum majus* L., *Petunia hybrida* Vilm., and *Arabidopsis thaliana* (L.) Heynh. (Winkel-Shirley, 2001; Tanaka et al., 2008; Dixon and Steele, 1999; Holton and Cornish, 1995). Generally, the following six enzymes are involved in the biosynthesis pathway of anthocyanin: chalcone synthase (CHS), chalcone isomerase (CHI), flavanone 3-hydroxylase (F3H), dihydroflavonol 4-reductase (DFR), anthocyanidin synthase

(ANS), and UDP-glucose-flavonoid 3-O-glucosyltransferase (UGFT) (Ma et al., 2009). In addition to structural genes, regulatory genes have been reported to play a vital role in the biosynthesis of anthocyanins in strawberry plants (Streisfeld and Rausher, 2010). Mutations in structural genes and transcription factor genes can affect the biosynthesis of anthocyanin (Streisfeld and Rausher, 2010).

Among structural genes, DFR encodes the first enzyme for anthocyanin biosynthesis in the flavonoid pathway (Griesbach, 2005). DFR controls the direction of carbon flux and catalyzes the conversion of dihydroflavones to unstable leucoanthocyanidins, which are common precursors for the anthocyanin and proanthocyanidin biosynthesis (Wang et al., 2013). The regulation of *DFR* expression has been well documented to change the flower color of Japanese parsley, lotus, and pansy (Hasegawa et al., 2001; Buathong et al., 2013; Li et al., 2014). Moreover, deactivation of DFR has been reported to result in loss of anthocyanins and proanthocyanidin in barley and *Arabidopsis* mutants (Olsen et al., 1993; Shirley et al., 1995).

Fragaria pentaphylla Losinsk is a wild diploid species of the *Fragaria* genus that belongs to the Rosaceae family (Yu, 1974). The

* Corresponding author.

E-mail address: lijmtzc@126.com (J. Li).

two different fruit colors (white and red) can be found in *F. pentaphylla* varieties in the field (Yu, 1974). The genome of *F. pentaphylla* is small and is considered a good model for studies of the mechanisms that underlie the production of fruit color. In the present study, we compared the expression of six structural genes and one regulatory gene, which are involved in the biosynthesis pathway of anthocyanin in white and red fruits of *F. pentaphylla*. To determine whether *FpDFR* is involved in the anthocyanin biosynthesis in *F. pentaphylla*, we examined the relationship between the *FpDFR* gene and the contents of both anthocyanin and flavonoids, isolated the cDNA and full-length sequences of the *FpDFR* gene, and analyzed the associations between single nucleotide polymorphism (SNPs) and fruit color. The results of this study provide a basic reference for the regulation of fruit color in strawberry.

2. Materials and methods

2.1. Plant materials and samples collection

Nine individual plants with non-matured white fruits and nine individual plants with red fruits of the diploid *F. pentaphylla* were collected respectively in July 2012 in Mao County, Chengdu City, Sichuan Province, China. The plants were transplanted and grown in a walk-in growth chamber (Taizhou University, Zhejiang Province, China), at day/night temperature of 20 °C/15 °C for 10 h/14 h. When the fruits were matured, two fruits and one young leaf were collected from every plant. The tissues were immediately frozen by liquid nitrogen and stored at –80 °C until analysis.

2.2. Quantitative RT-PCR

Real-time quantitative PCR (qPCR) was performed on the cDNA libraries of three red fruit and three white fruit respectively using specific primers (*CHS*: FP5' CCGACTACTACT TTCGT ATCACC A 3', RP5' ACTACCACCATGTCTGTCTTGC 3'; *CHI*: FP5' TGACAATGATACTACCGCTGAC 3', RP5' CTGTGGGAAGGTCTGATCTTT3'; *F3H*: FP5' TGGAGAGATGTGACAAAGCAGT 3', RP5' TCAAATGCCTCTTCTCTAAACC3'; *DFR*: FP 5' CCAAGGACCCTGA-GAATGAA 3', RP 5' TTACTCTCCGGCAAATTCG 3'; *ANS*: FP5' TGTGGCAACAAGTGAGTATGC3', RP5' ACCGACCTCCTTCTC-CAGCCT3'; *UFGT*: FP5' AATGGCATGCTTAAGAGTTTGA3', RP5' CTGTTGTGCGAGTTGTTTAGT 3'; *MYB10*: FP5' AGATGCAGGAA-GAGCTGTAGA C 3', RP5' TTCCTAGAGCTTATGAAGCCT 3') with the SYBR Green Master Mix (SYBR® Premix Ex Taq™, TaKaRa, Co. Ltd., Dalian, China) on an Illumina Eco Real-Time PCR System (Illumina Inc., San Diego, CA, USA). Reactions were performed in triplicate using 12.5 μL of 10 × buffer, 0.4 mol L⁻¹ of each primer, 2 μL of diluted cDNA, and nuclease-free water to a total volume of 20 μL. The real-time qPCR assay conditions were as follows: pre-incubation at 95 °C for 30 s, then 40 cycles of 95 °C for 5 s, 60 °C for 30 s, and final extension at 72 °C for 30 s. The length of the PCR product was not less than 100 bp. Fluorescence was measured at the end of each annealing step. Results from the real-time qPCR assays were normalized using the Ct value corresponding to the reference gene (18S rRNA) using specific primers (FP 5' ACCGTTGATTCGCACAATTGGTCATCG 3' and RP 5' TACTGCGGGTGGCAATCGGACG 3'). Relative expression levels were quantified using the 2^{-ΔΔ} method (Pfaffl, 2001).

2.3. Determination of total anthocyanin content

Three red fruits and three white fruits, which were collected from the same plants used for cDNA cloning, were separately ground and extracted in 15 mL of methanol acidified with 1.0% HCl at room temperature for 24 h. The extract was centrifuged at

12000×g for 5 min, and the supernatant was transferred to a new centrifuge tube. The total anthocyanin concentration was determined using a modified pH differential method described previously (Garzon and Wrolstad, 2002). A general spectrophotometer (T6 New Century; Purkinje General Instrument Co. Ltd., Beijing, China) was used to measure the absorbance of the extract solution in buffers of pH 1.0 and pH 4.5 at 520 nm and 700 nm, respectively. Total anthocyanins content was expressed as mg of cyanidin-3-glucoside per g of dry weight of fruit. Total anthocyanins = $A \times MW \times B \times 100 \times V/\xi$, where $A = [(A_{520} - A_{700})_{pH\ 1.0} - (A_{520} - A_{700})_{pH\ 4.5}]$, MW indicates molecular weight 433.2 g mol⁻¹, B indicates the dilution ratio, V indicates the final volume, and ξ indicates the molar extinction coefficient of 22,400. All samples were analyzed in triplicate.

2.4. Determination of total flavonoid content

The total flavonoid content was determined using an aluminum chloride colorimetric assay (Jia et al., 1999). A specific volume of extract was placed in a 10-mL volumetric flask and diluted with distilled water to 5 mL. Then, 0.3 mL of NaNO₂ (1:20) and 3 mL of AlCl₃ (1:10) were added after 5 min, and after a further 6 min, 2 mL of 1 M NaOH were added. Then, distilled water was added to obtain a total volume of 10 mL. The solution was mixed again, and a general spectrophotometer (T6 New Century, Purkinje General Instrument Co., Beijing, China) was used to measure absorbance at 510 nm. Total flavonoid content was expressed as mg rutin equivalents per g of fresh mass. All samples were analyzed in triplicate.

2.5. Cloning and characteristics of *FpDFR* cDNA

Total RNA was extracted from red fruits, using the RNeasy Plant Mini Kit (Qiagen Co. Ltd., Hilden, Germany). The RNA concentration was quantified using a NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific Inc., Wilmington, DE, USA) and the absorbance ratios at 260 nm and 280 nm were calculated to evaluate the quality of RNA and to diluted RNA to 1 μg μl⁻¹. A cDNA library was synthesized using the PrimeScript™ Double Strand cDNA Synthesis Kit (Takara Co. Ltd., Dalian, China) with oligo dT primers.

The full-length coding sequence for the DFR protein was PCR amplified. For this, a cDNA library was used, which was amplified using total RNA extracted from a red fruit as template. PCR amplification was conducted with specific primers (FP 5' ATGGGATCG-GAGTCCGAATCC 3' and RP' TTAGCCAGTGACTTCGACATGG 3'), designed according to the potential cDNA region of the genomic DNA of *F. pentaphylla*, using the following PCR amplification procedure: pre-denaturation at 95 °C for 5 min, followed by 30 cycles of amplification (denaturation at 95 °C for 30 s, annealing at 50 °C for 30 s, and extension at 72 °C for 1 min), and a final extension at 72 °C for 10 min. PCR products were analyzed with 1.0% agarose gels, single fragments were recovered from gels, and purified using a DNA purification kit (BioTeke Co. Ltd., Beijing, China). The purified gene product was ligated into the pMD18-T vector (Takara Co. Ltd., Dalian, China) and transformed into *E. coli* DH 5α for subsequent sequencing. PCR products were analyzed on 1.0% agarose gels and recovered from the gel via DNA purification kit (BioTeke Co. Ltd., Beijing, China). PCR products were directly sequenced by Biosune Co. Ltd. (Shanghai, China) via the ABI 373X (ABI, Carlsbad, CA, USA). The nucleotide sequence of the *FpDFR* cDNA was analyzed via BLASTN search against the GenBank database of the National Center for Biotechnology Information (NCBI) at www.ncbi.nlm.nih.gov/BLAST/.

Both the theoretical isoelectric point and molecular weight were calculated online (http://www.expasy.ch/tools/pi_tool.html). The

ORF of the full-length DFR was identified via Genscan (<http://genes.mit.edu/GENSCAN.html>) and motifs were found with Plantcare (<http://bioinformatics.psb.ugent.be/webtools/plantcare/html/>). The secondary structure of the predicted DFR protein was constructed with the PredictProtein tool (<http://www.predictprotein.org/>) and the three-dimensional structure of the predicted DFR was modeled via SWISS-MODEL with Automated model (<http://swissmodel.expasy.org/>). A phylogenetic analysis of the FpDFR protein was performed based on the deduced protein sequence using the NJ method, which was implemented in MEGA 5.0.

2.6. Cloning of FpDFR genomic DNA and SNP analysis

Young leaves from *F. pentaphylla* with red fruits were collected and genomic DNA was extracted, following the protocol of the utilized DNA extraction kit (Dingguo Co. Ltd., Beijing, China). PCR amplification was conducted with specific primers (FP 5' GATCG-TATCAGACATTCTCC 3' and RP 5'AATGATGAATGAGACGTCCTA 3'), which were designed in accordance with the DFR gene sequence of *F. vesca* (GenBank accession No. NM_001305268.1). The following PCR conditions were utilized: pre-denaturation at 95 °C for 5 min, followed by 30 cycles of amplification (denaturation at 95 °C for 30 s, annealing at 55 °C for 30 s, and extension at 72 °C for 1 min), followed by a final extension at 72 °C for 10 min. Fragments were amplified using the Phusion® High-Fidelity DNA polymerase (New England Biolabs Inc., Ipswich, MA, USA). PCR products were

analyzed with 1.0% agarose gels and recovered from the gel via DNA purification kit (BioTeke Corporation, Beijing, China). PCR products were directly sequenced by Biosune Co. Ltd. (Shanghai, China) via the ABI 373X (ABI, Carlsbad, USA). The sequence was then mapped to determine the positions of SNPs. Significance levels of associations between SNPs and color trait were evaluated via SPSS software (version 19.0).

3. Results

3.1. Expression of genes involved in the anthocyanin biosynthesis pathway and their correlation with total anthocyanin and flavonoid contents in red and white fruits

Both the expression levels of structural genes (*FpCHS*, *FpDFR*, *FpANS*, and *FpUFGT*) and the regulatory gene *MYB10*, which is involved in the pathway of anthocyanin biosynthesis in red fruits were significantly higher than those in white fruits, while the expression of the remaining structural genes (*FpCHI* and *FpF3H*) were similar between red fruits and white fruits (Fig. 1).

The total anthocyanin content of red fruits was 12.36 ± 0.12 mg/100 g FW, while that of white fruits was almost undetectable (0.06 ± 0.08 mg/100 g FW) (Fig. 2a) and was approximately 200-fold lower than that of red fruits of *F. pentaphylla*. However, the total flavonoid content of red *F. pentaphylla* fruits was similar to that of white fruits (Fig. 2b).

3.2. Correlation between SNPs of FpDFR and fruit color

The full length of the *FpDFR* gene sequence of red *F. pentaphylla* fruits was 2080 bp. This sequence was submitted to GenBank under the accession number KR075887.1. A comparison of the full-length genomic sequence of *FpDFR* with that of the cDNA in the red *F. pentaphylla* fruits revealed six exons of 118, 170, 195, 160, 193, and 217 bp, separated by five introns of 91, 495, 108, 84, and 79 bp, respectively. It shared 95% homology with the *DFR* gene of *F. vesca* (GenBank accession number AY017480).

Twenty-seven SNPs were detected in *FpDFR* gDNA sequences between red and white fruits of *F. pentaphylla* (Table 1). Among these, transition substitutions were more frequent than transversions (66.7% vs. 33.3%) and more nucleotide variants were located in introns than in exons (70.4% vs. 29.6%) (Table 1). Transition between G and A was more common than between C and T. For transversions, the substitutions were as follows (arranged in order of decreasing frequency): C and G, T and G, and C and A. Transversions between C and A were not detected. A Chi-square




Fig. 1. Log₂ fold-change of anthocyanin biosynthesis pathway genes expression. The results represent the relative expression in red fruits vs. white fruits of *Fragaria pentaphylla*. CHS: chalcone synthase; CHI: chalcone isomerase; F3H: flavanone 3-hydroxylase; DFR: dihydroflavonol 4-reductase; ANS: anthocyanidin synthase; UFGT: UDP-glucose-flavonoid 3-O-glucosyltransferase.




Fig. 2. The total anthocyanin content (a) and total flavonoid content (b) in red (R) and white (W) fruits of *Fragaria pentaphylla*. Error bars indicate standard deviations of three independent biological samples.

Table 1
SNPs in *DFR* gene between red and white fruits and their association with fruit color in *Fragaria pentaphylla*.

Position	Location	Allel	Frequency		Chi-square (p value)	Type of single-base substitution
			Red fruit	White fruit		
43	Exon	C	0.444	0.889	2.492 (0.114)	Transversion
		G	0.556	0.111		
161	Intron	C	0.444	0.111	2.492 (0.114)	Transversion
		G	0.556	0.889		
248	Exon	A	0.556	1.000	6.923 (0.009)	Transition
		G	0.444	0.000		
362	Exon	C	0.222	0.222	0.000 (1.000)	Transversion
		G	0.778	0.778		
468	Intron	A	1.000	0.556	5.143(0.023)	Transition
		G	0.000	0.444		
471	Intron	A	0.000	0.222	2.250 (0.134)	Transition
		G	1.000	0.778		
596	Intron	C	0.000	0.222	2.250 (0.134)	Transversion
		G	1.000	0.778		
686	Intron	A	0.778	1.000	2.250 (0.134)	Transition
		G	0.222	0.000		
694	Intron	C	0.667	0.889	1.286 (0.257)	Transition
		T	0.333	0.111		
712	Intron	C	1.000	0.778	2.250 (0.134)	Transition
		T	0.000	0.222		
715	Intron	C	1.000	0.778	2.250 (0.134)	Transversion
		G	0.000	0.222		
736	Intron	A	1.000	0.778	2.250 (0.134)	Transition
		G	0.000	0.222		
775	Intron	C	0.000	0.222	2.250 (0.134)	Transversion
		G	1.000	0.778		
794	Intron	G	1.000	0.778	2.250 (0.134)	Transversion
		T	0.000	0.222		
808	Intron	G	0.000	0.222	2.250 (0.134)	Transversion
		T	1.000	0.778		
830	Intron	A	0.000	0.333	3.60 (0.058)	Transversion
		T	1.000	0.667		
1029	Intron	C	0.778	1.000	2.250 (0.134)	Transition
		T	0.222	0.000		
1340	Intron	A	0.222	0.000	2.250 (0.134)	Transition
		G	0.778	1.000		
1349	Intron	C	1.000	0.778	2.250 (0.134)	Transition
		T	0.000	0.222		
1381	Intron	C	0.778	1.000	2.250 (0.134)	Transition
		T	0.222	0.000		
1402	Exon	A	0.000	0.444	5.143 (0.023)	Transition
		G	1.000	0.556		
1450	Exon	C	1.000	0.778	2.250 (0.134)	Transition
		T	0.000	0.222		
1621	Intron	A	0.000	0.222	2.250 (0.134)	Transition
		G	1.000	0.778		
1656	Exon	A	0.222	0.000	2.250 (0.134)	Transition
		G	0.778	1.000		
1668	Exon	A	0.222	0.222	0.000 (1.000)	Transition
		G	0.778	0.778		
1703	Exon	A	0.000	0.333	3.60 (0.058)	Transition
		G	1.000	0.667		
1908	Intron	C	0.778	1.000	2.250 (0.134)	Transition
		T	0.222	0.000		

test showed that only three SNPs significantly associated with fruit color (Table 1). The SNP at 248 bp was classified as a nonsense mutation, resulting in an amino acid substitution of lysine for glutamic acid in white fruits of *F. pentaphylla*, while SNPs at positions 468 and 1402 bp were classified as synonymous mutations.

3.3. Characterization of *FpDFR* cDNA and proteins

The full-length cDNA of *FpDFR* in red fruits of *F. pentaphylla* had a length of 1050 bp, encoding a polypeptide of approximately 350 amino acid residues, which corresponds to a molecular mass of 38.992 kDa, with a theoretical pI of 6.2 (Fig. 3a; sequence was submitted to GenBank under accession number KR075887). The nucleotide sequence of *FpDFR* cDNA in red fruits of *F. pentaphylla*

shared significant homologies with the *DFR* cDNA of *Fragaria vesca* L. cultivar *Alexandria* (KC894050.1, 99% similarity), *Fragaria* × *ananassa* Duch. cultivar *Elsanta* (KC894054.1, 99% similarity), *Fragaria* × *ananassa* cultivar *Chandler* (AF029685.1, 99% similarity), *Rosa rugosa* Thunb. (KM203111.1, 92% similarity), a *Rosa* hybrid cultivar (AB490072.1, 91% similarity), and *Rosa chinensis* Jacquem. (KF734592.1, 90% similarity).

An analysis of the secondary structure of the predicted *FpDFR* protein revealed that all examined *FpDFRs* contained an NAD (P)-binding domain. Furthermore, secondary structure analysis of the predicted *FpDFR* revealed that this predominantly consisted of β -pleated sheets (50.29%), followed by α -helices (36.29%), and random coils (13.43%) (Fig. 3b).

The quality of the predicted three-dimensional structure of the




Fig. 3. The full-length cDNA sequence of the *FpDFR* gene and its deduced amino acid sequence (a), three-dimensional structure (b) and secondary structure (c). The arrows indicate α -helices, rectangles indicate β -pleated sheets, and gray indicates the NAD (P)-binding domain.

FpDFR protein in red fruits of *F. pentaphylla* was revealed via comparison with data in the Protein Data Base Structures database, and the resulting global model quality estimate was 0.83, suggesting high quality of the predicted model (Fig. 3c).

The phylogenetic tree based on the nucleic acid sequences of *FpDFR* in red fruits of *F. pentaphylla* and other nucleic acid sequences of *DFR* of various plant species are shown in Fig. 4. Analysis showed the *FpDFR* sequence in red fruits of *F. pentaphylla* to be closely related to the *DFR* nucleic acid sequence of *F. vesca* (Fig. 4).

4. Discussion

Here, the detected expression levels of four structural genes (*FpCHS*, *FpDFR*, *FpANS*, and *FpUFGT*) and the regulatory gene *FpMYB10* in red fruits of *F. pentaphylla* were significantly higher compared to white fruits, while anthocyanin contents followed an identical pattern. These results indicated that expression of these structural and regulatory genes was involved in the determination of *F. pentaphylla* fruit color. Mutations in regulatory genes (*MYB* in particular) are always expected to have fewer deleterious pleiotropic effects than mutations in structural genes due to high tissue

specificity of the transcription factors (Streisfeld and Rausher, 2009). For example, a nonsense mutation in the *MYB1* protein may be responsible for the apparent color difference between white and red fruits of the Chinese bayberry (Niu et al., 2010). An independent insertion of a retrotransposon into the transcription factor *Ruby* has been reported to result in the red color of the blood orange (Butelli et al., 2012). Although these examples highlight several cases supporting this prediction, the possibility that mutations in structural genes were also involved cannot be ruled out (Durbin et al., 2003; Zufall and Rausher, 2004). The involvement of structural genes and regulatory genes in the biosynthesis of fruit anthocyanins is well documented in some plant species, such as *Capsicum* spp., *Prunus avium* (Wei et al., 2015), and *Vaccinium myrtillus* (Jaakola et al., 2003). However, to the best of our knowledge, our results form the first demonstration of associations between gene expression and fruit color in *F. pentaphylla*.

DFR is a ventral gene involved in anthocyanin biosynthesis. It has been isolated in many species, such as *Z. mays* (Meyer et al., 1992; Schwarz-Sommer et al., 1987), *Bromheadia finlaysoniana* Lindl. (Liew et al., 1998), *P. hybrida* (Chu et al., 2014), *Gerbera hybrida* Hort. (Helariutta et al., 1993), *F. × ananassa* (Moyano et al., 1998),




Fig. 4. Phylogenetic analysis of the *FpDFR* cDNA sequence (in gray) and those of other plant species generated using the neighbor-joining method. In the phylogenetic tree, species names and GenBank accession numbers are provided in parentheses. Numbers at branch nodes refer to bootstrap values (100 replicates).

and *F. vesca* (Shulaev et al., 2011). However, the function of DFR to determine plant fruit color varies among plant species (Wang et al., 2004; Ju et al., 1997). Wang et al. (2013) reported that DFR gene expression has a significant relationship with the anthocyanin content in the purple sweet potato. However, Wang et al. (2004) reported that DFR gene expression has no relationship with the anthocyanin content in the pericarp of Litchi. Ju et al. (1997) found that 'Delicious,' 'Golden delicious,' and 'Indo' apples also contained relatively high DFR activity, although they did not accumulate anthocyanins. Salvatierra et al. (2013) reported that DFR gene expression in two botanical forms of *Fragaria chiloensis* Duch. spp. *chiloensis* was only equal in red and white botanical forms during the ripening stages. In our study, the strong correlation between the expression of the *FpDFR* gene and the anthocyanin content in *F. pentaphylla* indicates *FpDFR* expression to be a key step in the anthocyanin biosynthesis. Lin et al. (2013) verified via RNAi technology, that DFR is a key gene in the pathway of anthocyanin biosynthesis in *F. x ananassa* fruits.

SNPs have been revealed as the most common source of genetic variation in eukaryotic species and consequently, have become an important marker for genetic studies, such as identifying candidate genes that affect diverse phenotypes (Wondji et al., 2007). In this study, we found that exon and intron sequences showed high similarity and only 27 SNPs were detected in a genomic DNA *FpDFR* gene comparison between red and white fruits of *F. pentaphylla*. Transition substitutions were more common than transversions (with a ratio of transitions to transversions close to 2:1), which was also observed for *Drosophila* (Moriyama and Powell, 1996), the mosquito *Anopheles funestus* (Wondji et al., 2007), and humans (Brookes, 1999).

Among all SNPs, only three significantly associated with fruit color. SNPs at positions 468 and 1402 bp were classified as synonymous mutations. Sauna and Kimchi-Sarfaty (2011) found that synonymous SNPs could have significant phenotypic effects, influencing the conformation, function, and expression of proteins. The regulatory effects of synonymous SNPs were verified to be strongly associated with CpG islands (Karambataki et al., 2014).

However, we did not detect such a CpG island. In addition, the higher frequency of C ↔ T and A ↔ G SNPs transitions was likely, at least in part, related to 5-methylcytosine deamination reactions that frequently occur at CpG dinucleotides in particular (Holliday and Grigg, 1993; Wondji et al., 2007). The SNP at 248 bp was classified as a nonsense mutation, resulting in an amino acid substitution of lysine for glutamic acid in white fruits of *F. pentaphylla*. However, the substitution did not change the secondary mRNA structure (Appendix 1) nor the tertiary protein structure (Appendix 2). In addition, based on the secondary structures of the *FpDFR* protein in red fruits of *F. pentaphylla*, mutated amino acids were not located in key structural regions, such as the α -helix, β -sheet, or substrate binding sites. Therefore, this SNP is a candidate genetic marker for red color; however, it would be classified as a non-conservative amino acid substitution and might not have a major impact on protein function.

An alternative mechanism might better explain color control in *F. pentaphylla*, i.e., trans-regulation. The coordinated expression of multiple structural genes in fruits of *F. pentaphylla* further suggests a trans-mutation in *FpDFR* (similar to MYB mutations), instead of a cis-mutation of *FpDFR*, regulating anthocyanin biosynthesis in *F. pentaphylla*. Lin-Wang et al. (2014) reported that MYB10 has the ability to auto-regulate itself as well as regulate all enzymes throughout the biosynthesis pathway of *F. vesca*, from the first committed step (CHS) to the final step (UFGT). Similar results have been reported for *Citrus sinensis* Osbeck (Lo Piero et al., 2006; Lo Piero, 2015). Further studies about the trans-regulation of anthocyanin biosynthesis are required to elucidate the mechanisms that underlie the different fruit colors of *F. pentaphylla*.

Acknowledgement

We thank Wanqing Yang for the kind help with plant collection. We would also like to thank the two anonymous reviewers for helpful comments. This work was financially supported by the National Natural Science Foundation of China (No. 31261120580).

Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.bse.2017.04.004>.

References

- Brookes, A.J., 1999. The essence of SNPs. *Gene* 234 (2), 177–186.
- Buathong, R., Saetiew, K., Phansiri, S., Parinthewong, N., Arunyanart, S., 2013. Tissue culture and transformation of the antisense *DFR* gene into lotus (*Nelumbo nucifera* Gaertn.) through particle bombardment. *Sci. Hortic.* 161 (24), 216–222.
- Butelli, E., Licciardello, C., Zhang, Y., Liu, J.J., Mackay, S., Bailey, P., Reforgiato-Rsecupero, G., Martina, C., 2012. Retrotransposons control fruit-specific, cold-dependent accumulation of anthocyanins in blood oranges. *Plant Cell* 24, 1242–1255.
- Chu, Y.X., Pan, J.S., Wu, A.Z., Cai, R., Chen, H.R., 2014. Molecular cloning and functional characterization of dihydroflavonol-4-reductase gene from *Calibrachoa hybrida*. *Sci. Hortic.* 165, 398–403.
- Dixon, R.A., Steele, C.L., 1999. Flavonoids and isoflavonoids—a gold mine for metabolic engineering. *Trends Plant Sci.* 4, 394–400.
- Durbin, M.L., Lundy, K.E., Morrell, P.L., Torres-Martinez, C.L., Clegg, M.T., 2003. Genes that determine flower color: the role of regulatory changes in the evolution of phenotypic adaptations. *Mol. Phylogenet. Evol.* 29, 507–518.
- Gasiorowski, K., Szyba, K., Brokos, B., Iaczyńska, B.K., Iodarczyk, M.J., Oszmiański, J., 1997. Antimutagenic activity of anthocyanins isolated from *Aronia melanocarpa* fruits. *Cancer Lett.* 119 (1), 37–46.
- Garzon, G.A., Wrolstad, R.E., 2002. Comparison of the stability of pelargonidin-based anthocyanins in strawberry juice and concentrate. *J. Food Sci.* 67 (4), 1288–1299.
- Griesbach, R.J., 2005. Biochemistry and genetics of flower color. *Plant Breed. Rev.* 25, 89–114.
- Goto, T., Kondo, T., 1991. Structure and molecular stacking of anthocyanins—flower color variation. *Angew. Chem. Int. Ed. Engl.* 30, 17–33.
- Hasegawa, H., Fukasawa-Akada, T., Okuno, T., Niizeki, M., Suzuki, M., 2001. Anthocyanin accumulation and related gene expression in Japanese parsley (*Oenanthe stolonifera*, DC.) induced by low temperature. *J. Plant Physiol.* 158, 71–78.
- Helariutta, Y., Elomaa, P., Kotilainen, M., Seppanen, P., Teeri, T.H., 1993. Cloning of cDNA coding for dihydroflavonol-4-reductase (DFR) and characterization of *dfr* expression in the corollas of *Gerbera hybrida* var. Regina (Compositae). *Plant Mol. Biol.* 22 (2), 183–193.
- Holton, T.A., Cornish, E.C., 1995. Genetics and biochemistry of anthocyanin biosynthesis. *Plant Cell* 7, 1071–1083.
- Holliday, R., Grigg, G.W., 1993. DNA methylation and mutation. *Mutat. Res.* 285 (1), 61–67.
- Jaakola, L., Hohtola, A., Määttä, K., Törrönen, S., Kärenlampi, S., 2003. Flavonoid biosynthesis in bilberry (*Vaccinium myrtillus* L.). *Acta Hortic.* 618, 415–419.
- Jia, Z.S., Tang, M.C., Wu, J.M., 1999. The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chem.* 64 (4), 555–559.
- Ju, Z.G., Yuang, Y.B., Liu, C.L., Wang, Y.Z., Tian, X.P., 1997. Dihydroflavonol reductase activity and anthocyanin accumulation in 'Delicious', 'Golden delicious' and 'Indo' apples. *Sci. Hortic.* 70 (1), 31–43.
- Kähkönen, M.P., Heinonen, M., 2003. Antioxidant activity of anthocyanins and their aglycons. *J. Agr. Food Chem.* 51 (3), 628–633.
- Karambataki, M., Malousi, A., Kouidou, S., 2014. Risk-associated coding synonymous SNPs in type 2 diabetes and neurodegenerative diseases: genetic silence and the underrated association with splicing regulation and epigenetics. *Mutat. Res.* 770, 85–93.
- Koide, T., Kamei, H., Hashimoto, Y., Kojima, T., Hasegawa, M., 1996. Antitumor effect of hydrolyzed anthocyanin from grape rinds and red rice. *Cancer Biother. Radio* 11 (4), 273–277.
- Li, Q., Wang, J., Sun, H.Y., Shang, X., 2014. Flower color patterning in pansy (*Viola × wittrockiana* Gams.) is caused by the differential expression of three genes from the anthocyanin pathway in acyanic and cyanic flower areas. *Plant Physiol. Bioch* 84, 134–141.
- Liew, C.F., Loh, C.S., Gho, C.J., Lim, S.H., 1998. The isolation, molecular characterization and expression of dihydroflavonol 4-reductase cDNA in the orchid *Bromheadia finlaysonian*. *Plant Sci.* 135, 161–169.
- Lin, X., Xiao, M., Luo, Y., Wang, J.Y., Wang, H.Q., 2013. The effect of RNAi-induced silencing of *FaDFR* on anthocyanin metabolism in strawberry (*Fragaria × ananassa*) fruit. *Sci. Hortic.* 160, 123–128.
- Lin-Wang, K., McGhie, T.K., Wang, M., Liu, Y.H., Warren, B., Storey, R., Easley, R.V., Allan, A.C., 2014. Engineering the anthocyanin regulatory complex of strawberry (*Fragaria vesca*). *Front. Plant Sci.* 5, 1–14.
- Lo Piero, A.R., Puglisi, I., Petrone, G., 2006. Gene characterization, analysis of expression and in vitro synthesis of dihydroflavonol 4-reductase from *Citrus sinensis* (L.) Osbeck. *Phytochemistry* 67, 684–695.
- Lo Piero, A.R., 2015. The State of the art in biosynthesis of anthocyanins and its regulation in pigmented sweet oranges [(*Citrus sinensis*) L. Osbeck]. *J. Agr. Food Chem.* 63, 4031–4041.
- Ma, H.M., Pooler, M., Griesbach, R., 2009. Anthocyanin regulatory/structural gene expression in *Phalaenopsis*. *J. Am. Soc. Hortic. Sci.* 134, 88–96.
- Meyer, P., Linn, F., Heidmann, I., Meyer, H., Niedenhof, I., Saedler, H., 1992. Endogenous and environmental factors influence 35S promoter methylation of a maize A1 gene construct in transgenic petunia and its colour phenotype. *Mol. Gen. Genet.* 231 (3), 345–352.
- Moriyama, E.N., Powell, J.R., 1996. Intraspecific nuclear DNA variation in *Drosophila*. *Mol. Biol. Evol.* 13 (1), 261–277.
- Moyano, E., Portero-Robles, I., Medina-Escobar, N., Valpuesta, V., Muñoz-Blanco, J., Caballero, J.L., 1998. A fruit-specific putative dihydroflavonol 4-reductase gene is differentially expressed in strawberry during the ripening process. *Plant Physiol.* 117 (2), 711–716.
- Niu, S.S., Xu, C.J., Zhang, W.S., Zhang, B., Li, X., Wang, K.L., Ferguson, I.B., Allan, A.C., Chen, K.S., 2010. Coordinated regulation of anthocyanin biosynthesis in Chinese bayberry (*Myrica rubra*) fruit by a R2R3 MYB transcription factor. *Planta* 231, 887–899.
- Olsen, O., Wang, X., VonWettstein, D., 1993. Sodium azide mutagenesis: preferential generation of A-T → G-C transitions in the barley Ant18 gene. *P. Natl. Acad. Sci. U. S. A.* 90, 8043–8047.
- Pfaffl, M.W., 2001. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* 29, 2002–2007.
- Salvatierra, A., Pimentel, P., Moya-León, M.A., Herre, R., 2013. Increased accumulation of anthocyanins in *Fragaria chiloensis* fruits by transient suppression of *FcMYB1* gene. *Phytochemistry* 90, 25–36.
- Sauna, Z.E., Kimchi-Sarfaty, C., 2011. Understanding the contribution of synonymous mutations to human disease. *Nat. Rev. Genet.* 12, 683–691.
- Schwarz-Sommer, Z., Shepherd, N., Tacke, E., Gierl, A., Rohde, W., Leclercq, L., Mattes, M., Berndtgen, R., Peterson, P.A., Saedler, H., 1987. Influence of transposable elements on the structure and function of the A1 gene of *Zea mays*. *EMBO J.* 6 (2), 287–294.
- Shirley, B.W., Kubasek, W.L., Storz, G., Bruggemann, E., Koornneef, M., Ausubel, F.M., Goodman, H.M., 1995. Analysis of *Arabidopsis* mutants deficient in flavonoid biosynthesis. *Plant J.* 8, 659–671.
- Shulaev, V., Sargent, D.J., Crowhurst, R.N., Mockler, T.C., Folkerts, O., Delcher, A.L., Jaiswal, P., Mockett, K., Liston, A., Mane, S.P., et al., 2011. The genome of woodland strawberry (*Fragaria vesca*). *Nat. Genet.* 43, 109–116.
- Streisfeld, M.A., Rauscher, M.D., 2009. Altered trans-regulatory control of gene expression in multiple anthocyanin genes contributes to adaptive flower color evolution in *Mimulus aurantiacus*. *Mol. Biol. Evol.* 26 (2), 433–444.
- Streisfeld, M.A., Rauscher, M.D., 2010. Population genetics, pleiotropy, and the preferential fixation of mutations during adaptive evolution. *Evolution* 65, 629–642.
- Tanaka, Y., Sasaki, N., Ohmiya, A., 2008. Biosynthesis of plant pigments: anthocyanins, betalains and carotenoids. *Plant J.* 54 (4), 733–749.
- Wang, H.B., Nair, M.G., Strasburg, G.M., Chang, Y.C., Booren, A.M., Gray, J.L., DeWitt, D.L., 1999. Antioxidant and anti-inflammatory activities of anthocyanins and their aglycon, cyanidin, from tart cherries. *J. Nat. Prod.* 62, 294–296.
- Wang, H.X., Fan, W.J., Li, H., Yang, J., Huang, J.R., Zhang, P., 2013. Functional characterization of dihydroflavonol-4-reductase in anthocyanin biosynthesis of purple sweet potato underlies the direct evidence of anthocyanins function against abiotic stresses. *PLoS ONE* 8 (11), e78484.
- Wang, H.Z., Huang, X.M., Hu, G.B., Huang, H.B., 2004. Studies on the relationship between anthocyanin biosynthesis and related enzymes in litchi pericarp. *Agr. Sci. China* 37, 2028–2032.
- Wei, H.R., Chen, X., Zong, X.J., Shu, H.R., Gao, D.S., Liu, Q.Z., 2015. Comparative transcriptome analysis of genes involved in anthocyanin biosynthesis in the red and yellow fruits of sweet cherry (*Prunus avium* L.). *PLoS ONE* 10 (3), e0121164.
- Winkel-Shirley, B., 2001. Flavonoid biosynthesis. A colorful model for genetics, biochemistry, cell biology, and biotechnology. *Plant Physiol.* 126, 485–493.
- Wondji, C.S., Hemingway, J., Ranson, H., 2007. Identification and analysis of single nucleotide polymorphisms (SNPs) in the mosquito *Anopheles funestus*, malaria vector. *BMC Genomics* 8, 5–18.
- Yu, D.J., 1974. Rosaceae. In: Flora of China, vol 37. Science Press, Beijing.
- Zufall, R.A., Rauscher, M.D., 2004. Genetic changes associated with floral adaptation restrict future evolutionary potential. *Nature* 428, 847–850.